arXiv:hep-ph/9901221v1 5 Jan 1999

Charged Higgs Associated Production with W at Linear Collider

Shou Hua Zhu

CCAST (World Laboratory), P. O. Box 8730, Beijing, 100080, P. R. China

Institute of Theoretical Physics, Academia Sinica, P.O. Box 2735, Beijing 100080, P.R. China

ABSTRACT

We have calculated the cross section of the process $e^+e^- \to W^\pm H^\mp$ at the linear collider in the supersymmetrical two-higgs-doublet model. We find that the cross section decreases rapidly with the increment of $\tan\beta$, and which can reach almost 1 fb for the favorable parameters.

PACS number(s): 12.15.Lk, 12.60.Jv, 14.80.Cp

Keyword(s): Higgs Physics, Two-Higgs-doublet Model, Linear Collider

I. INTRODUCTION

The Higgs boson is the missing piece and also the least known one of the standard model (SM) and beyond. Under the framework of the minimal supersymmetrical standard model (MSSM) [1], there contain charged Higgs bosons, which are one of the most distinguished signal compared with SM. To find this kind of charged Higgs boson and study its properties are one of the primary goals of the present and next generation of colliders.

The production of the charged Higgs boson at hadron colliders can mainly via top quark decay if kinetically allowed; recently, we have studied single charged Higgs production through bg channel and found this channel could be helpful in finding the charged Higgs [2]. Otherwise, the heavy charged Higgs would be very hard to produce at hadron colliders. At linear collider, the primary mechanism to produce charged Higgs bosons are $e^+e^- \to H^+H^-$ [3] or $\gamma\gamma \to H^+H^-$ [4]. However, the production rates for both processes drop rapidly with the increment of the charged Higgs mass. As a complementary channel in discovering charged Higgs, although suffering from the low rate (this channel is the one loop process), $e^+e^- \to W^\pm H^\mp$ channel is important in studying the $\gamma - W^\pm - H^\mp$ and $Z - W^\pm - H^\mp$ vertexes. In fact, these two vertexes have been studied through the decay processes of the charged Higgs boson [5].

In literatures, there are many works on neutral Higgs associated production with gauge bosons (γ, Z and W) both on hadron and linear colliders [6], which are potentially useful in dealing with the backgrounds. However, the charged Higgs bosons associated production with W at linear collider is deserved a study, which will be the topic of this paper. At linear collider, there is an alternative collision mode: $\gamma\gamma$ collision, which is commonly thought the most suitable place to study the Higgs properties. The $W^\pm H^\mp$ pair production in this collision mode will be presented in the forthcoming work [7].

II. FORMALISM OF CHARGED HIGGS BOSON PRODUCTION ASSOCIATED WITH W

The Feynman diagrams of the process $e^+(p_1)e^-(p_2) \rightarrow H^\pm(k_1)W^\mp(k_2)$ are shown in Fig 1, and the corresponding Feynman rules could be found in Ref. [8]. We perform the calculation in the 't Hooft-Feynman gauge and use dimensional regularization to all the ultraviolet divergence in the virtual loop contributions.

As usual, we define the mandelstam variables as

$$S = (p_1 + p_2)^2 = (k_1 + k_2)^2$$

$$T = (p_1 - k_1)^2 = (p_2 - k_2)^2$$

$$U = (p_1 - k_2)^2 = (p_2 - k_1)^2. \tag{1}$$

The amplitude of the process can be written as

$$M = \bar{U}(p_2)\left(f_1 \not{\epsilon} P_R + f_2 \not{\epsilon} P_L + \not{k}_2\left((f_3 P_R + f_4 P_L) p_1.\epsilon + (f_5 P_R + f_6 P_L) p_2.\epsilon\right)\right) V(p_1), \tag{2}$$

where ϵ is the polarization vector of W boson and $P_{L,R} = (1 \mp \gamma_5)/2$. The form factors f_i could be expressed as

$$f_i = \sum_{j=1}^{11} f_i^{(j)}, \tag{3}$$

where j corresponds to the contributions from the diagrams index in Fig. 1. For simplicity, if possible, the form factors $f_i^{(j)}$ are divided into $f_i^{(j),\gamma}$ and $f_i^{(j),Z}$, which are from the s-channel γ and Z contributions, respectively; for diagram (2) in Fig. 1, the form factors are further divided into fermionic and bosonic contributions. The non-vanishing form factors are explicitly presented in Appendix.

The cross section of process $e^+(p_1)e^-(p_2) \rightarrow H^\pm(k_1)W^\mp(k_2)$ can be written as

$$\sigma = \int_{T_{min}}^{T_{max}} \frac{1}{16\pi S^2} \overline{\sum_{spins}} |M|^2 \, dT \tag{4}$$

with

$$T_{min} = \frac{1}{2}(m_{H^\pm}^2 + m_W^2 - S - \sqrt{(S - (m_{H^\pm} + m_W)^2)(S - (m_{H^\pm} - m_W)^2)})$$

$$T_{max} = \frac{1}{2}(m_{H^\pm}^2 + m_W^2 - S + \sqrt{(S - (m_{H^\pm} + m_W)^2)(S - (m_{H^\pm} - m_W)^2)}) \qquad (5)$$

where the bar over the summation recalls average over initial fermions spins.

III. NUMERICAL RESULTS AND DISCUSSIONS

In the following we present some numerical results of charged Higgs boson associated production with W in the process of $e^+e^- \to H^\pm W^\mp$. In our calculations, we choose $m_W = 80.33GeV$, $m_Z = 91.187GeV$, $m_t = 176GeV$, $m_b = 4.5GeV$ and $\alpha = 1/128$. For simplicity, we choose the parameters of the Higgs sectors which satisfied the MSSM relations, i.e., there are two independent parameters $\tan\beta$ and $m_{H^\pm}$ [8].

In Fig. 2, the $W^\pm H^\mp$ pair production cross sections as a function of the $m_{H^\pm}$ at linear collider are presented, where $\sqrt{S} = 500GeV$. From the figures, we can see that the cross sections drops rapidly with the increment of $\tan\beta$, and for $\tan\beta = 2$ and $m_{H^\pm} = 100 \sim 250$ GeV, the cross sections are insensitive to the mass of the charged Higgs boson, which is around 0.2 fb. We note that the peak around 180 GeV is due to the threshold effect that the decay channel of the charged Higgs boson to top and bottom are opened.

In Fig. 3, the $W^\pm H^\mp$ pair production cross sections as a function of the $\sqrt{S}$ are presented for $m_{H^\pm} = 200$ GeV. We can see that there is a peak around 400 GeV, which is the consequence of the competitive between the enlargement of phase space and the s-channel suppression. For three cases of $\tan\beta$, the cross sections of small $\tan\beta$ are larger than that of large $\tan\beta$, which can reach 1 fb for favorable parameters.

To summarize, we have calculated the cross section of the process $e^+e^- \to W^\pm H^\mp$ at the next linear collider in the supersymmetrical two-higgs-doublet model. We find that the cross section decreases rapidly with the increment of $\tan\beta$, and which can reach almost 1

fb for the favorable parameters. We note that in supersymmetrical framework including MSSM, the s-particles can also contribute to this process, which will be studied in the near future.

ACKNOWLEDGMENTS

This work was supported in part by the post doctoral foundation of China and the author gratefully acknowledges the support of K.C. Wong Education Foundation, Hong Kong.

---

[1] H.P. Nilles, Phys. Rep. **110**, 1 (1984); H.E. Haber and G.L. Kane, Phys. Rep. **117**, 75 (1985).

[2] C.S. Huang and S.H. Zhu, submitted to Phys. Rev. Lett., hep-ph/9812201.

[3] A. Djouadi, J. Kalinowski and P.M. Zerwas, Z. Phys. **C70**, 435 (1996); A. Arhrib, M.C. Peyranere and G. Moultaka, Phys. Lett. **B341**, 313 (1995).

[4] D. Bowser-Chao, K. cheung and S. Thomas, Phys. Lett **B315**, 399 (1983); W.G. Ma, C.S. Li and L. Han, Phys. Rev. **D53**, 1304 (1996); Erratum-ibid, Phys. Rev. **D56**, 4420 (1997); S.H. Zhu, C.S. Li and C.S. Gao, Phys. Rev. **D58**, 05007 (1998).

[5] M. Peyranere, H.E. Haber and P. Irulegui, Phys. Rev. D **44**, 191 (1991); M. Mendez and A. Pomarol, Nucl. Phys. **B349**, 369 (1991); S. Kanemura, hep-ph/9710237;

[6] See for examples, A. Barroso, J. Pulido and J.C. Romao, Nucl. Phys. **B267**, 509 (1986); A. Abbasabadi, D. Bowser-Chao, D.A. Dicus and W.W. Repko, Phys. Rev. **D52**, 3919 (1995);ibid, Phys. Rev. **D57**, 550 (1998); ibid, Phys. Rev. **D58**, 057301 (1998); A. Djouadi, V. Driesen, W. Hollik and J. Rosiek, Nucl. Phys. **B491**, 68 (1997).

[7] S.H. Zhu, in preparation.

[8] J.F. Gunion et. al., Higgs Hunter's Guide (Addison-Wesley, MA,1990).

[9] G. Passarino and M. Veltman, Nucl.Phys.B**33** (1979)151; A. Denner, Fortschr. Phys. 41(1993)4; R.Mertig, M.Bohm and A.Denner, Comp.Phys.Comm.**64** (1991)345.



FIG. 1. The typical Feynman diagrams for the process of $e^+e^- \rightarrow H^\pm W^\mp$.



FIG. 2. The $W^{\pm}H^{\mp}$ pair production cross sections as a function of the $m_{H^{\pm}}$ at linear collider, where $\sqrt{S} = 500GeV$, and the solid, dashed and dotted lines represent $\tan\beta = 2$, 10 and 40, respectively.



FIG. 3. The $W^{\pm}H^{\mp}$ pair production cross sections as a function of the $\sqrt{S}$ at linear collider, where $m_{H^{\pm}} = 200 GeV$, and the solid, dashed and dotted lines represent $\tan\beta = 2$, 10 and 40, respectively.

APPENDIX

In this Appendix, the non-zero form factors defined in Eq. 3 are explicitly written as

$$f_2^{(1)} = \frac{\alpha^2 m_W}{4\sin^4\theta_W}\cos(\alpha-\beta)\sin(\alpha-\beta)\Big[(T-m_H^2)D_0(1)-(T-m_{h_0}^2)D_0(2)$$
$$+(T-m_{H^\pm}^2)\left(D_1(2)-D_1(1)\right)+(T-m_W^2)\left(D_3(1)-D_3(2)\right)\Big], \quad (6)$$

$$f_6^{(1)} = \frac{\alpha^2 m_W}{\sin^4\theta_W}\cos(\alpha-\beta)\sin(\alpha-\beta)\left(D_3(1)-D_3(2)\right), \quad (7)$$

$$f_1^{(2),\gamma,F} = \frac{\alpha^2}{m_W\sin^2\theta_W S}\Big[-m_t^2\cot\beta B_0(S,m_b^2,m_b^2)+2m_b^2\tan\beta B_0(S,m_t^2,m_t^2)$$
$$+m_t^2(m_b^2-m_t^2-T)\cot\beta C_0(1)+2m_b^2(m_b^2-m_t^2+U)\tan\beta C_0(2)$$
$$+2\Big[(m_t^2\cot\beta(U-m_{H^\pm}^2)+m_b^2\tan\beta(U+m_{H^\pm}^2)\Big]C_1(2)$$
$$-\Big[(m_t^2\cot\beta(T+m_{H^\pm}^2)+m_b^2\tan\beta(T-m_{H^\pm}^2)\Big]C_1(1)$$
$$-2\Big[(m_t^2\cot\beta(T-m_W^2)-m_b^2\tan\beta(U+m_W^2)\Big]C_2(2)$$
$$-\Big[(m_t^2\cot\beta(T+m_W^2)+m_b^2\tan\beta(m_W^2-U)\Big]C_2(1)$$
$$-2(m_t^2\cot\beta+m_b^2\tan\beta)(2C_{00}(2)-C_{00}(1))\Big], \quad (8)$$

$$f_2^{(2),\gamma,F} = f_1^{(2),\gamma,F}(T\leftrightarrow U), \quad (9)$$

$$f_3^{(2),\gamma,F} = \frac{2\alpha^2}{m_W\sin^2\theta_W S}\Big[2m_b^2\tan\beta C_0(2)$$
$$+2(2m_b^2\tan\beta+m_t^2\cot\beta)C_1(2)-m_t^2\cot\beta C_1(1)$$
$$+m_b^2\tan\beta(2C_2(2)+C_2(1))+(m_b^2\tan\beta+m_t^2\cot\beta)(2C_{11}(2)-C_{11}(1))$$
$$+(m_b^2\tan\beta+m_t^2\cot\beta)(2C_{12}(2)-C_{12}(1))\Big], \quad (10)$$

$$
\begin{aligned}
f_4^{(2),\gamma,F} = \frac{2\alpha^2}{m_W \sin^2\theta_W S}\Big[& -m_t^2\cot\beta C_0(1) \\
& -(2m_t^2\cot\beta + m_b^2\tan\beta)C_1(1) + 2m_b^2\tan\beta C_1(2) \\
& -m_t^2\cot\beta(2C_2(2)+C_2(1)) + (m_b^2\tan\beta + m_t^2\cot\beta)(2C_{11}(2)-C_{11}(1)) \\
& +(m_b^2\tan\beta + m_t^2\cot\beta)(2C_{12}(2)-C_{12}(1))\Big],
\end{aligned}
\tag{11}
$$

$$
f_5^{(2),\gamma,F} = f_4^{(2),\gamma,F}, \tag{12}
$$

$$
f_6^{(2),\gamma,F} = f_3^{(2),\gamma,F}, \tag{13}
$$

$$
\begin{aligned}
f_1^{(2),Z,F} = \frac{\alpha^2}{2\sin^2\theta_W\cos^2\theta_W m_W(s-m_Z^2)} & \Big[\Big((3-2\sin^2\theta_W)m_t^2\cot\beta + 3m_b^2\tan\beta\Big) B_0(S,m_b^2,m_b^2) \\
& -\Big((3-4\sin^2\theta_W)m_b^2\tan\beta + 3m_t^2\cot\beta\Big) B_0(S,m_t^2,m_t^2) \\
& +m_t^2\Big((T+m_t^2)\cot\beta(3-2\sin^2\theta_W) + 2m_b^2\sin^2\theta_W\cot\beta + 3m_b^2\tan\beta\Big) C_0(2) \\
& -m_b^2\Big((U+m_b^2)\tan\beta(3-4\sin^2\theta_W) + 4m_t^2\sin^2\theta_W\tan\beta + 3m_t^2\cot\beta\Big) C_0(1) \\
& +\Big(\cot\beta m_t^2\Big(-4m_{H^\pm}^2\sin^2\theta_W - (3-4\sin^2\theta_W)U\Big) - \tan\beta m_b^2(3-4\sin^2\theta_W)(m_{H^\pm}^2+U)\Big) C_1(2) \\
& -\Big(\tan\beta m_b^2\Big(-2m_{H^\pm}^2\sin^2\theta_W + (3-2\sin^2\theta_W)T\Big) - \cot\beta m_t^2(3-2\sin^2\theta_W)(m_{H^\pm}^2+T)\Big) C_1(1) \\
& +\Big(\cot\beta m_t^2\Big(-4T\sin^2\theta_W - (3-4\sin^2\theta_W)m_W^2\Big) - \tan\beta m_b^2(3-4\sin^2\theta_W)(m_W^2+U)\Big) C_2(2) \\
& +\Big(\tan\beta m_b^2\Big(2U\sin^2\theta_W + (3-2\sin^2\theta_W)m_W^2\Big) + \cot\beta m_t^2(3-2\sin^2\theta_W)(m_W^2+T)\Big) C_2(1) \\
& +2(m_t^2\cot\beta + m_b^2\tan\beta)\Big((3-4\sin^2\theta_W)C_{00}(2) - (3-2\sin^2\theta_W)C_{00}(1)\Big)\Big],
\end{aligned}
\tag{14}
$$

$$
f_2^{(2),Z,F} = \frac{-1+2\sin^2\theta_W}{2\sin^2\theta_W} f_1^{(2),Z,F}(T\leftrightarrow U), \tag{15}
$$

$$f_3^{(2),Z,F} = \frac{\alpha^2}{m_W \sin^2\theta_W \cos^2\theta_W (S - m_Z^2)} \Big[m_b^2 \tan\beta(-3 + 4\sin^2\theta_W) C_0(2)$$

$$+(2m_b^2 \tan\beta + m_t^2 \cot\beta)(-3 + 4\sin^2\theta_W) C_1(2) - m_t^2 \cot\beta(-3 + 2\sin^2\theta_W) C_1(1)$$

$$+m_b^2 \tan\beta((-3 + 4\sin^2\theta_W) C_2(2) + 2C_2(1)) + (m_b^2 \tan\beta + m_t^2 \cot\beta) \times$$

$$\Big((-3 + 4\sin^2\theta_W) C_{11}(2) - (-3 + 2\sin^2\theta_W) C_{11}(1)\Big)$$

$$+(m_b^2 \tan\beta + m_t^2 \cot\beta)((-3 + 4\sin^2\theta_W) C_{12}(2) - (-3 + 2\sin^2\theta_W) C_{12}(1))\Big], \quad (16)$$

$$f_4^{(2),Z,F} = \frac{\alpha^2(-1 + 2\sin^2\theta_W)}{2m_W \cos^2\theta_W \sin^2\theta_W S} \Big[m_t^2 \cot\beta(3 - 2\sin^2\theta_W) C_0(1)$$

$$+(3 - 2\sin^2\theta_W)(2m_t^2 \cot\beta + m_b^2 \tan\beta) C_1(1) + m_b^2 \tan\beta(-3 + 4\sin^2\theta_W) C_1(2)$$

$$-m_t^2 \cot\beta(4C_2(2) + (-3 + 2\sin^2\theta_W) C_2(1))$$

$$+(m_b^2 \tan\beta + m_t^2 \cot\beta)((-3 + 4\sin^2\theta_W) C_{11}(2) - (-3 + 2\sin^2\theta_W) C_{11}(1))$$

$$+(m_b^2 \tan\beta + m_t^2 \cot\beta)((-3 + 4\sin^2\theta_W) C_{12}(2) - (-3 + 2\sin^2\theta_W) C_{12}(1))\Big], \quad (17)$$

$$f_5^{(2),Z,F} = \frac{2\sin^2\theta_W}{-1 + 2\sin^2\theta_W} f_4^{(2),Z,F}, \tag{18}$$

$$f_6^{(2),Z,F} = \frac{-1 + 2\sin^2\theta_W}{2\sin^2\theta_W} f_3^{(2),Z,F}, \tag{19}$$

$$f_1^{(2),\gamma,B} = \frac{\alpha^2 m_W}{2S \sin^2\theta_W} \Big[-\cos(\alpha - \beta) \Big((S + m_H^2)\sin(\alpha - \beta) + m_Z^2 \cos(\alpha + \beta)\sin(2\beta)\Big) C_0(3)$$

$$\sin(\alpha - \beta) \Big((S + m_{h_0}^2)\cos(\alpha - \beta) + m_Z^2 \sin(\alpha + \beta)\sin(2\beta)\Big) C_0(4)$$

$$+ \cos(\alpha - \beta)\sin(\alpha - \beta) \Big((2m_{H^\pm}^2 - T - U)(C_1(3) - C_1(4))$$

$$-(2m_W^2 - T - U)(C_2(3) - C_2(4))\Big)$$

$$+2\sin(\alpha-\beta)\left(-2\cos(\alpha-\beta)+\sec^2\theta_W\cos(\alpha+\beta)\cos(2\beta)\right)C_{00}(5)$$

$$+2\cos(\alpha-\beta)\left(2\sin(\alpha-\beta)+\sec^2\theta_W\cos(\alpha+\beta)\sin(2\beta)\right)C_{00}(3)$$

$$+2\cos(\alpha-\beta)\left(2\sin(\alpha-\beta)-\sec^2\theta_W\sin(\alpha+\beta)\cos(2\beta)\right)C_{00}(6)$$

$$+2\sin(\alpha-\beta)\left(-2\cos(\alpha-\beta)+\sec^2\theta_W\sin(\alpha+\beta)\sin(2\beta)\right)C_{00}(4)\Big], \tag{20}$$

$$f_2^{(2),\gamma,B}=f_1^{(2),\gamma,B}, \tag{21}$$

$$f_3^{(2),\gamma,B}=\frac{\alpha^2 m_W}{S\sin^2\theta_W}\Big[\sin(\alpha-\beta)\left(2\cos(\alpha-\beta)-\sec^2\theta_W\cos(2\beta)\cos(\alpha+\beta)\right)(C_1(5)+C_{11}(5)+C_{12}(5))$$

$$-\cos(\alpha-\beta)\cos(\alpha+\beta)\sin(2\beta)\sec^2\theta_W C_1(3)$$

$$+\cos(\alpha-\beta)\left(-2\sin(\alpha-\beta)+\sec^2\theta_W\cos(2\beta)\sin(\alpha+\beta)\right)(C_1(6)+C_{11}(6)+C_{12}(6))$$

$$-\sin(\alpha-\beta)\sin(\alpha+\beta)\sin(2\beta)\sec^2\theta_W C_1(4)$$

$$+2\cos(\alpha-\beta)\sin(\alpha-\beta)\left(C_2(3)-C_2(4)\right)$$

$$-\cos(\alpha-\beta)\left(2\sin(\alpha-\beta)+\sec^2\theta_W\cos(\alpha+\beta)\sin(2\beta)\right)(C_{11}(3)+C_{12}(3))$$

$$+\sin(\alpha-\beta)\left(2\cos(\alpha-\beta)-\sec^2\theta_W\sin(\alpha+\beta)\sin(2\beta)\right)(C_{11}(4)+C_{12}(4))\Big], \tag{22}$$

$$f_4^{(2),\gamma,B}=f_5^{(2),\gamma,B}=f_6^{(2),\gamma,B}=f_3^{(2),\gamma,B}, \tag{23}$$

$$f_1^{(2),Z,B}=\frac{\alpha^2 m_W}{2\sin^2\theta_W(S-m_Z^2)}\Big[\frac{1}{\cos^2\theta_W}\cos(\alpha-\beta)\sin(\alpha-\beta)\left(-B_0(S,m_H^2,m_Z^2)+B_0(S,m_{h_0}^2,m_Z^2)\right)$$

$$-\frac{\cos(\alpha-\beta)}{\cos^2\theta_W}\left(\sin(\alpha-\beta)(m_Z^2+2T+2U)+m_Z^2\tan^2\theta_W\cos(\alpha+\beta)\sin(2\beta)\right)C_0(13)$$

$$+\frac{\sin(\alpha-\beta)}{\cos^2\theta_W}\left(\cos(\alpha-\beta)(m_Z^2+2T+2U)-m_Z^2\tan^2\theta_W\sin(\alpha+\beta)\sin(2\beta)\right)C_0(14)$$

$$
\begin{aligned}
&-\cos(\alpha-\beta)\Big(\sin(\alpha-\beta)(m_Z^2-S-m_H^2)+m_Z^2\tan^2\theta_W\cos(\alpha+\beta)\sin(2\beta)\Big)\,C_0(3)\\
&+\sin(\alpha-\beta)\Big(\cos(\alpha-\beta)(m_Z^2-S-m_{h_0}^2)-m_Z^2\tan^2\theta_W\sin(\alpha+\beta)\sin(2\beta)\Big)\,C_0(4)\\
&+(-2m_{H^\pm}^2+T+U)\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_1(3)-C_1(4))\\
&+\frac{1}{\cos^2\theta_W}(2m_{H^\pm}^2+T+U)\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_1(14)-C_1(13))\\
&+(-2m_W^2+T+U)\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_2(4)-C_2(3))\\
&+\frac{1}{\cos^2\theta_W}(2m_W^2+T+U)\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_2(14)-C_2(13))\\
&+\frac{(\cos^2\theta_W-\sin^2\theta_W)}{\cos^2\theta_W}\sin(\alpha-\beta)\Big(-2\cos(\alpha-\beta)+\sec^2\theta_W\cos(\alpha+\beta)\cos(2\beta)\Big)\,C_{00}(5)\\
&-\frac{(\cos^2\theta_W-\sin^2\theta_W)\cos(\alpha-\beta)}{\cos^2\theta_W\sin^2\theta_W}\Big(2\sin(\alpha-\beta)+\sec^2\theta_W\cos(\alpha+\beta)\sin(2\beta)\Big)\,C_{00}(3)\\
&+\frac{\sin(\alpha-\beta)}{\cos^4\theta_W}\Big(-2\sec^2\theta_W\cos(\alpha-\beta)+\cos(2\beta)\cos(\alpha+\beta)\Big)\,C_{00}(7)\\
&-\frac{\cos(\alpha-\beta)}{\cos^4\theta_W}\Big(-2\sec^2\theta_W\sin(\alpha-\beta)+\cos(2\beta)\sin(\alpha+\beta)\Big)\,C_{00}(8)\\
&-\frac{(\sin^2\theta_W-\cos^2\theta_W)}{\cos^4\theta_W}\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_{00}(9)-C_{00}(10))\\
&+\frac{(\cos^2\theta_W-\sin^2\theta_W)}{\cos^2\theta_W}\cos(\alpha-\beta)\Big(-2\sin(\alpha-\beta)+\sec^2\theta_W\sin(\alpha+\beta)\cos(2\beta)\Big)\,C_{00}(6)\\
&-\frac{(\cos^2\theta_W-\sin^2\theta_W)\sin(\alpha-\beta)}{\cos^4\theta_W}\Big(2\cos^2\theta_W\cos(\alpha-\beta)-\sin(\alpha+\beta)\sin(2\beta)\Big)\,C_{00}(4)\\
&-\frac{2}{\cos^2\theta_W}\cos(\alpha-\beta)\sin(\alpha-\beta)\,(C_{00}(11)-C_{00}(12))\\
&+\frac{\cos(\alpha-\beta)}{\cos^2\theta_W}\Big(2\sin(\alpha-\beta)+\sec^2\theta_W\sin(2\beta)\cos(\alpha+\beta)\Big)\,C_{00}(13)\\
&+\frac{\sin(\alpha-\beta)}{\cos^2\theta_W}\Big(-2\cos(\alpha-\beta)+\sec^2\theta_W\sin(2\beta)\sin(\alpha+\beta)\Big)\,C_{00}(14),
\end{aligned}
\tag{24}
$$

$$f_2^{(2),Z,B} = \frac{-1 + 2\sin^2\theta_W}{2\sin^2\theta_W} f_1^{(2),Z,B}, \tag{25}$$

$$\begin{aligned}
f_3^{(2),Z,B} = \frac{\alpha^2 m_W}{2\sin^2\theta_W (S - m_Z^2)} \Big[& \frac{2}{\cos^2\theta_W} \cos(\alpha-\beta)\sin(\alpha-\beta) \Big(C_0(11) - C_0(12) + \sec^2\theta_W \left(C_0(14) - C_0(13)\right) \\
& + \frac{(\cos^2\theta_W - \sin^2\theta_W)}{\cos^4\theta_W} \sin(\alpha-\beta) \Big(-2\cos^2\theta_W \cos(\alpha-\beta) + \cos(2\beta)\cos(\alpha+\beta) \Big) \times \\
& (C_1(5) + C_{11}(5) + C_{12}(5)) + \frac{(\cos^2\theta_W - \sin^2\theta_W)}{\cos^4\theta_W} \cos(\alpha-\beta)\cos(\alpha+\beta)\sin(2\beta) C_1(3) \\
& + \frac{1}{\cos^4\theta_W} \sin(\alpha-\beta) \Big(2\cos^2\theta_W \cos(\alpha-\beta) - \cos(2\beta)\cos(\alpha+\beta) \Big) (C_1(7) + C_{11}(7) + C_{12}(7)) \\
& + \frac{1}{\cos^4\theta_W} \cos(\alpha-\beta) \Big(-2\cos^2\theta_W \sin(\alpha-\beta) + \cos(2\beta)\sin(\alpha+\beta) \Big) (C_1(8) + C_{11}(8) + C_{12}(8)) \\
& + \frac{(\cos^2\theta_W - \sin^2\theta_W)}{\cos^4\theta_W} \cos(\alpha-\beta)\sin(\alpha-\beta) (C_1(9) - C_1(10) - C_{11}(9) + C_{11}(10) - C_{12}(9) \\
& + C_{12}(10)) + \frac{\cos^2\theta_W - \sin^2\theta_W}{\cos^4\theta_W} \cos(\alpha-\beta) \Big(2\cos^2\theta_W \sin(\alpha-\beta) - \cos(2\beta)\sin(\alpha+\beta) \Big) \\
& (C_1(6) + C_{11}(6) + C_{12}(6)) + \frac{\cos^2\theta_W - \sin^2\theta_W}{\cos^4\theta_W} \sin(\alpha-\beta)\sin(\alpha+\beta)\sin(2\beta) C_1(4) \\
& + \frac{2}{\cos^2\theta_W} \sin(\alpha-\beta)\cos(\alpha-\beta) (2C_1(11) - 2C_1(12) + C_{11}(11) - C_{11}(11) + C_{12}(11) - C_{12}(11)) \\
& - \frac{1}{\cos^4\theta_W} \cos(\alpha-\beta) \Big(4\cos^2\theta_W \sin(\alpha-\beta) + \cos(\alpha+\beta)\sin(2\beta) \Big) C_1(13) \\
& - \frac{1}{\cos^4\theta_W} \sin(\alpha-\beta) \Big(-4\cos^2\theta_W \cos(\alpha-\beta) + \sin(\alpha+\beta)\sin(2\beta) \Big) C_1(14) \\
& + 4\cos(\alpha-\beta)\sin(\alpha-\beta) (C_2(4) - C_2(3)) \\
& + \frac{2}{\cos^2\theta_W} \cos(\alpha-\beta)\sin(\alpha-\beta) (C_2(11) - C_2(12) + C_2(13) - C_2(14)) \\
& + \frac{(\cos^2\theta_W - \sin^2\theta_W)}{\cos^4\theta_W} \cos(\alpha-\beta) \Big(\cos(\alpha+\beta)\sin(2\beta) + 2\cos^2\theta_W \sin(\alpha+\beta) \Big) (C_{11}(3) + C_{12}(3))
\end{aligned}$$

$$
\begin{aligned}
&+\frac{\cos^2\theta_W-\sin^2\theta_W}{\cos^4\theta_W}\sin(\alpha-\beta)\left(\sin(\alpha+\beta)\sin(2\beta)-2\cos^2\theta_W\sin(\alpha-\beta)\right)\left(C_{11}(4)+C_{12}(4)\right)\\
&-\frac{1}{\cos^4\theta_W}\cos(\alpha-\beta)\left(2\cos^2\theta_W\sin(\alpha-\beta)+\cos(\alpha+\beta)\sin(2\beta)\right)\left(C_{11}(13)+C_{12}(13)\right)\\
&-\frac{1}{\cos^4\theta_W}\sin(\alpha-\beta)\left(-2\cos^2\theta_W\cos(\alpha-\beta)+\sin(\alpha+\beta)\sin(2\beta)\right)\left(C_{11}(14)+C_{12}(14)\right)\Big],
\end{aligned}
\tag*{(2}
$$

$$
f_4^{(2),Z,B}=f_6^{(2),Z,B}=\frac{-1+2\sin^2\theta_W}{2\sin^2\theta_W}f_3^{(2),Z,B},
\tag{27}
$$

$$
f_5^{(2),Z,B}=f_3^{(2),Z,B},
\tag{28}
$$

$$
\begin{aligned}
f_1^{(3),\gamma}=&\frac{\alpha^2}{4\sin^2\theta_W\cos^2\theta_W(m_W^2-m_{H^\pm}^2)m_W S}\Pi_{G^\pm H^\pm}\\
&-\frac{\alpha^2(m_W^2-S)}{4\sin^2\theta_W\cos\theta_W(m_W^2-m_{H^\pm}^2)m_W m_{H^\pm}^2 S}\Pi_{W^\pm H^\pm},
\end{aligned}
\tag{29}
$$

$$
\begin{aligned}
f_1^{(3),Z}=&\frac{\alpha^2 m_Z}{4\cos^3\theta_W(m_W^2-m_{H^\pm}^2)m_W^2(S-m_Z^2)}\Pi_{G^\pm H^\pm}\\
&+\frac{\alpha^2(m_W^2-S)}{4\sin^2\theta_W\cos\theta_W(m_W^2-m_{H^\pm}^2)m_W m_{H^\pm}^2(S-m_Z^2)}\Pi_{W^\pm H^\pm},
\end{aligned}
\tag{30}
$$

$$
\begin{aligned}
f_1^{(4),\gamma}=&\frac{\alpha^2\left(m_W m_Z\sin(2\beta)\sin(\alpha+\beta)-\cos\theta_W S\cos(\alpha-\beta)\right)}{8\sin^2\theta_W\cos^2\theta_W m_{h_0}^2 m_W(m_W^2-m_{H^\pm}^2)S}\Pi_{h_0}^T\\
&-\frac{\alpha^2\left(m_W m_Z\sin(2\beta)\cos(\alpha+\beta)+\cos\theta_W S\sin(\alpha-\beta)\right)}{8\sin^2\theta_W\cos^2\theta_W m_H^2 m_W(m_W^2-m_{H^\pm}^2)S}\Pi_H^T,
\end{aligned}
\tag{31}
$$

$$
\begin{aligned}
f_1^{(4),Z}=&\frac{\alpha^2\left(\cos(\alpha-\beta)\cos^2\theta_W(m_Z^2-S)-m_Z^2\sin^2\theta_W\sin(2\beta)\sin(\alpha+\beta)\right)}{8\sin^2\theta_W\cos^3\theta_W m_{h_0}^2 m_W(m_W^2-m_{H^\pm}^2)(m_Z^2-S)}\Pi_{h_0}^T\\
&+\frac{\alpha^2\left(\sin(\alpha-\beta)\cos^2\theta_W(m_Z^2-S)+m_Z^2\sin^2\theta_W\sin(2\beta)\cos(\alpha+\beta)\right)}{8\sin^2\theta_W\cos^3\theta_W m_H^2 m_W(m_W^2-m_{H^\pm}^2)(m_Z^2-S)}\Pi_H^T,
\end{aligned}
\tag{32}
$$

$$
f_1^{(5),\gamma}=-\frac{\alpha^2\cos(\alpha-\beta)}{8\sin^2\theta_W\cos\theta_W m_{h_0}^2 m_W S}\Pi_{h_0}^T-\frac{\alpha^2\sin(\alpha-\beta)}{8\sin^2\theta_W\cos\theta_W m_H^2 m_W S}\Pi_H^T,
\tag{33}
$$

$$f_1^{(5),Z} = \frac{\alpha^2 \sin^2\theta_W \cos(\alpha-\beta)}{8\cos^3\theta_W m_{h_0}^2 m_W(+m_Z^2 - S)}\Pi^T_{h_0} + \frac{\alpha^2\sin^2\theta_W\sin(\alpha-\beta)}{8\cos^3\theta_W m_H^2 m_W(m_Z^2-S)}\Pi^T_H, \tag{34}$$

$$\begin{aligned} f_1^{(6),\gamma} = \frac{\alpha^2 m_Z}{4\sin^2\theta_W\cos\theta_W S}\Big[&\sin(\alpha-\beta)\left(2\sec^2\theta_W\cos(\alpha-\beta) - \cos(2\beta)\cos(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_H^2, m_{H^\pm}^2) \\ &-\cos(\alpha-\beta)\left(\sec^2\theta_W\sin(\alpha-\beta) + \sin(2\beta)\cos(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_H^2, m_W^2) \\ &+\cos(\alpha-\beta)\left(-2\sec^2\theta_W\sin(\alpha-\beta) + \cos(2\beta)\sin(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_{H^\pm}^2, m_{h_0}^2) \\ &+\sin(\alpha-\beta)\left(\sec^2\theta_W\cos(\alpha-\beta) - \sin(2\beta)\sin(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_{h_0}^2, m_W^2)\Big], \end{aligned} \tag{35}$$

$$\begin{aligned} f_1^{(6),Z} = \frac{\alpha^2 m_Z}{4\cos^3\theta_W(S-m_Z^2)}\Big[&\sin(\alpha-\beta)\left(2\sec^2\theta_W\cos(\alpha-\beta) - \cos(2\beta)\cos(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_H^2, m_{H^\pm}^2) \\ &-\cos(\alpha-\beta)\left(\sec^2\theta_W\sin(\alpha-\beta) + \sin(2\beta)\cos(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_H^2, m_W^2) \\ &+\cos(\alpha-\beta)\left(-2\sec^2\theta_W\sin(\alpha-\beta) + \cos(2\beta)\sin(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_{H^\pm}^2, m_{h_0}^2) \\ &+\sin(\alpha-\beta)\left(\sec^2\theta_W\cos(\alpha-\beta) - \sin(2\beta)\sin(\alpha+\beta)\right)B_0(m_{H^\pm}^2, m_{h_0}^2, m_W^2)\Big], \end{aligned} \tag{36}$$

$$f_1^{(7),\gamma} = \frac{\alpha^2 m_W}{2\sin^2\theta_W S}\cos(\alpha-\beta)\sin(\alpha-\beta)\left(B_0(m_W^2, m_H^2, m_W^2) - B_0(m_W^2, m_{h_0}^2, m_W^2)\right), \tag{37}$$

$$f_1^{(7),Z} = \frac{\alpha^2 m_W}{2\cos^2\theta_W(S-m_Z^2)}\cos(\alpha-\beta)\sin(\alpha-\beta)\left(B_0(m_W^2, m_H^2, m_W^2) - B_0(m_W^2, m_{h_0}^2, m_W^2)\right), \tag{38}$$

$$f_1^{(8),Z} = \frac{\alpha^2 m_Z}{2\cos^3\theta_W(S-m_Z^2)}\cos(\alpha-\beta)\sin(\alpha-\beta)\left(B_0(S, m_H^2, m_Z^2) - B_0(S, m_{h_0}^2, m_Z^2)\right), \tag{39}$$

$$\begin{aligned} f_1^{(9),\gamma} = \frac{\alpha^2 m_W}{8\cos^2\theta_W\sin^2\theta_W(m_{H^\pm}^2 - m_W^2)S}\Big[&\cos(2\beta)\sin(2\beta)\left(A_0(m_{A_0}^2) + 4A_0(m_{H^\pm}^2) - A_0(m_Z^2) - 4A_0(m_W^2)\right) \\ &-\left(\cos^2\theta_W\cos(2\beta)\sin(2\alpha) + \sin^2\theta_W\cos(2\alpha)\sin(2\beta)\right)\left(A_0(m_H^2) - A_0(m_{h_0}^2)\right)\Big], \end{aligned} \tag{40}$$

$$\begin{aligned} f_1^{(9),Z} = \frac{\alpha^2 m_W}{8\cos^4\theta_W(m_{H^\pm}^2 - m_W^2)(S-m_Z^2)}\Big[&\cos(2\beta)\sin(2\beta)\left(A_0(m_{A_0}^2) + 4A_0(m_{H^\pm}^2) - A_0(m_Z^2) - 4A_0(m_W^2)\right) \\ &-\left(\cos^2\theta_W\cos(2\beta)\sin(2\alpha) + \sin^2\theta_W\cos(2\alpha)\sin(2\beta)\right)\left(A_0(m_H^2) - A_0(m_{h_0}^2)\right)\Big], \end{aligned} \tag{41}$$

$$f_2^{(j),\gamma} = f_1^{(j),\gamma}, \text{ for } j = 3 - 9, \tag{42}$$

$$f_2^{(j),Z} = \frac{2\sin^2\theta_W}{-1 + 2\sin^2\theta_W} f_1^{(j),Z}, \text{ for } j = 3 - 9, \tag{43}$$

$$f_2^{(10)} = -\frac{\alpha^2}{8\sin\theta_W\cos\theta_W(m_W^2 - m_{H^\pm}^2)m_W m_{H^\pm}^2\pi^2}\Pi_{W^\pm H^\pm}, \tag{44}$$

$$\begin{aligned} f_2^{(11)} = & \frac{\alpha^2\cos(\alpha-\beta)}{16\sin^2\theta_W\cos\theta_W m_{h_0}^2 m_W(m_W^2 - m_{H^\pm}^2)}\Pi_{h_0}^T \\ & + \frac{\alpha^2\sin(\alpha-\beta)}{16\sin^3\theta_W\cos\theta_W m_H^2 m_W(m_W^2 - m_{H^\pm}^2)}\Pi_H^T, \end{aligned} \tag{45}$$

where

$$\begin{aligned} \Pi_{h_0}^T = & -24\cos\theta_W m_t^2\cos\alpha\csc\beta A_0(m_t^2) + 24\cos\theta_W m_b^2\sin\alpha\sec\beta A_0(m_b^2) \\ & - m_W m_Z\left(6\sin(\alpha-\beta) + \sin(\alpha+\beta)\cos(2\beta)\right)A_0(m_Z^2) \\ & - 2m_W\left(6\cos\theta_W m_W\sin(\alpha-\beta) + m_Z\cos(2\beta)\sin(\alpha+\beta)\right)A_0(m_W^2) \\ & - m_W m_Z\left(\cos(2\alpha)\sin(\alpha+\beta) + 2\cos(\alpha+\beta)\sin(2\alpha)\right)A_0(m_H^2) \\ & + 3m_W m_Z\cos(2\alpha)\sin(\alpha+\beta)A_0(m_{h_0}^2) + m_W m_Z\cos(2\beta)\sin(\alpha+\beta)A_0(m_{A_0}^2) \\ & + 2m_W\left(m_Z\cos(2\beta)\sin(\alpha+\beta) - 2\cos\theta_W m_W\sin(\alpha-\beta)\right)A_0(m_{H^\pm}^2), \end{aligned} \tag{46}$$

$$\begin{aligned} \Pi_H^T = & -24\cos\theta_W m_t^2\sin\alpha\csc\beta A_0(m_t^2) - 24\cos\theta_W m_b^2\cos\alpha\sec\beta A_0(m_b^2) \\ & + m_W m_Z\left(6\cos(\alpha-\beta) + \cos(\alpha+\beta)\cos(2\beta)\right)A_0(m_Z^2) \\ & + 2m_W\left(6\cos\theta_W m_W\cos(\alpha-\beta) + m_Z\cos(2\beta)\cos(\alpha+\beta)\right)A_0(m_W^2) \\ & + 3m_W m_Z\left(\cos(2\alpha)\cos(\alpha+\beta)\right)A_0(m_H^2) - m_W m_Z\cos(2\beta)\cos(\alpha+\beta)A_0(m_{A_0}^2) \end{aligned}$$

$$
\begin{aligned}
&+m_W m_Z \left(2\sin(2\alpha)\sin(\alpha+\beta)-\cos(2\alpha)\cos(\alpha+\beta)\right) A_0(m_{h_0}^2)\\
&-2m_W \left(m_Z\cos(2\beta)\cos(\alpha+\beta)-2\cos\theta_W m_W\cos(\alpha-\beta)\right) A_0(m_{H^\pm}^2),
\end{aligned}
\tag{47}
$$

$$
\begin{aligned}
\Pi_{G^\pm H^\pm} = {}& 12\cos^2\theta_W(m_t^2\cot\beta-m_b^2\tan\beta)A_0(m_t^2)\\
&+6\cos^2\theta_W(m_t^2-m_b^2)(m_b^2\tan\beta-m_t^2\cot\beta)B_0(0,m_b^2,m_t^2)\\
&+6\cos^2\theta_W\left(\cot\beta m_t^2(m_t^2-m_b^2-m_{H^\pm}^2)+\tan\beta m_b^2(m_t^2-m_b^2+m_{H^\pm}^2)\right)B_0(m_{H^\pm}^2,m_b^2,m_t^2)\\
&+\cos^2\theta_W(m_{h_0}^2-m_W^2)m_W^2\cos(\alpha-\beta)\sin(\alpha-\beta)B_0(0,m_{h_0}^2,m_W^2)\\
&-\cos^2\theta_W(m_H^2-m_W^2)m_W^2\cos(\alpha-\beta)\sin(\alpha-\beta)B_0(0,m_H^2,m_W^2)\\
&+m_W^2\left[2\cos\theta_W m_W\cos(\alpha-\beta)-m_Z\cos(2\beta)\cos(\alpha+\beta)\right]\left[\cos\theta_W m_W\sin(\alpha-\beta)\right.\\
&\left.+m_Z\sin(2\beta)\cos(\alpha+\beta)\right]B_0(m_{H^\pm}^2,m_H^2,m_{H^\pm}^2)\\
&+m_W^2\Big[m_Z^2\cos(2\beta)\cos^2(\alpha+\beta)\sin(2\beta)+\cos^2\theta_W\sin(\alpha-\beta)\Big(\\
&\cos(\alpha-\beta)(2m_H^2+2m_{H^\pm}^2-m_W^2)+m_Z^2\cos(2\beta)\cos(\alpha+\beta)\Big)\Big]B_0(m_{H^\pm}^2,m_H^2,m_W^2)\\
&+m_W^2\left[-2\cos\theta_W m_W\sin(\alpha-\beta)+m_Z\cos(2\beta)\sin(\alpha+\beta)\right]\left[\cos\theta_W m_W\cos(\alpha-\beta)\right.\\
&\left.-m_Z\sin(2\beta)\sin(\alpha+\beta)\right]B_0(m_{H^\pm}^2,m_{H^\pm}^2,m_{h_0}^2)\\
&+m_W^2\Big[m_Z^2\sin(2\beta)\cos(2\beta)\sin^2(\alpha+\beta)+\cos^2\theta_W\cos(\alpha-\beta)\Big(\\
&\sin(\alpha-\beta)(-2m_{h_0}^2-2m_{H^\pm}^2+m_W^2)-m_Z^2\sin(\alpha+\beta)\cos(2\beta)\Big)\Big]B_0(m_{H^\pm}^2,m_{h_0}^2,m_W^2),
\end{aligned}
\tag{48}
$$

$$
\begin{aligned}
\Pi_{W^\pm H^\pm} = {}& 6\cos\theta_W(m_b^2-m_t^2)(m_b^2\tan\beta+m_t^2\cot\beta)B_0(0,m_b^2,m_t^2)\\
&+6\cos\theta_W\left(\cot\beta m_t^2(m_t^2-m_b^2-m_{H^\pm}^2)+\tan\beta(m_b^2(m_t^2-m_b^2+m_{H^\pm}^2)\right)B_0(m_{H^\pm}^2,m_b^2,m_t^2)
\end{aligned}
$$

$$
\begin{aligned}
&+\sin(\alpha-\beta)m_W(m_{H^\pm}^2-m_H^2)\left(m_Z\cos(2\beta)\cos(\alpha+\beta)-2\cos\theta_W m_W\cos(\alpha-\beta)\right)\times\\
&\Big(B_0(0,m_H^2,m_{H^\pm}^2)-B_0(m_{H^\pm}^2,m_H^2,m_{H^\pm}^2)\Big)+\cos(\alpha-\beta)m_W(m_W^2-m_H^2)\times\\
&\left(m_Z\sin(2\beta)\cos(\alpha+\beta)+2\cos\theta_W m_W\sin(\alpha-\beta)\right)B_0(0,m_H^2,m_W^2)\\
&+m_W\cos(\alpha-\beta)\Big[\sin(\alpha-\beta)\cos\theta_W m_W(2m_H^2+3m_{H^\pm}^2-2m_W^2)\\
&+\cos(\alpha+\beta)\sin(2\beta)m_Z(m_H^2-m_W^2)\Big]B_0(m_{H^\pm}^2,m_H^2,m_W^2)\\
&-\cos(\alpha-\beta)m_W(m_{H^\pm}^2-m_{h_0}^2)\left(m_Z\cos(2\beta)\sin(\alpha+\beta)-2\cos\theta_W m_W\sin(\alpha-\beta)\right)\times\\
&\Big(B_0(0,m_{h_0}^2,m_{H^\pm}^2)-B_0(m_{H^\pm}^2,m_{h_0}^2,m_{H^\pm}^2)\Big)+\sin(\alpha-\beta)m_W(m_W^2-m_{h_0}^2)\times\\
&\left(m_Z\sin(2\beta)\sin(\alpha+\beta)-2\cos\theta_W m_W\cos(\alpha-\beta)\right)B_0(0,m_{h_0}^2,m_W^2)\\
&+m_W\sin(\alpha-\beta)\Big[\cos(\alpha-\beta)\cos\theta_W m_W(-2m_{h_0}^2-3m_{H^\pm}^2+2m_W^2)\\
&+\sin(\alpha+\beta)\sin(2\beta)m_Z(m_{h_0}^2-m_W^2)\Big]B_0(m_{H^\pm}^2,m_{h_0}^2,m_W^2). \qquad (49)
\end{aligned}
$$

In above equations, the definitions of B_0, C and D functions could be found in Ref. [9], the variables of C functions from 1 to 14 represent $(m_{H^\pm}^2, S, m_W^2, i)$, where i stand for the internal mass square of the loops, for $i = 1-14$, which are

$$
\begin{aligned}
&(m_t^2,m_b^2,m_b^2),(m_b^2,m_t^2,m_t^2),(m_H^2,m_W^2,m_W^2),(m_{h_0}^2,m_W^2,m_W^2)\\
&(m_H^2,m_{H^\pm}^2,m_{H^\pm}^2),(m_{h_0}^2,m_{H^\pm}^2,m_{H^\pm}^2),(m_{H^\pm}^2,m_H^2,m_{A_0}^2),(m_{H^\pm}^2,m_{h_0}^2,m_{A_0}^2),\\
&(m_{H^\pm}^2,m_Z^2,m_H^2),(m_{H^\pm}^2,m_Z^2,m_{h_0}^2),(m_W^2,m_{A_0}^2,m_H^2),(m_W^2,m_{A_0}^2,m_{h_0}^2),\\
&(m_W^2,m_H^2,m_Z^2),(m_W^2,m_{h_0}^2,m_Z^2), \qquad (50)
\end{aligned}
$$

and the variables of D functions from 1 to 2 represent $(m_{H^\pm}^2, 0, 0, m_W^2, T, S, i)$, where i stand for the internal mass square of the loops, for $i = 1-2$, which are

$$(m_H^2, m_W^2, 0, m_W^2), (m_{h_0}^2, m_W^2, 0, m_W^2). \tag{51}$$